



(How) Should ultrasound enter the match?

Orhan Güvener, MD¹, Yiğitcan Karanfil, MD², Levent Özçakar, MD³

¹Department of Physical and Rehabilitation Medicine, Mersin University Medical School, Mersin, Türkiye

²Hacettepe University Medical School, Sports Medicine, Ankara, Türkiye

³Department of Physical and Rehabilitation Medicine, Hacettepe University Medical School, Ankara, Türkiye

Owing to its several advantages, ultrasound (US) has gained importance in musculoskeletal medicine.^[1] Its applicability in sports arena/settings has also been established.^[2] Herein, we further extend its potential utility by asking “Why cannot it be used during the game as well?”. Similar to technologies such as video-assisted referee, “hawk eye” etc., why are physicians (or medical staff) unable to implement the play considering the athlete's health/injury? For instance, several degrees of acute lesions involving different tissues can readily be evaluated and further catastrophic injuries can be prevented. Indisputably, the medical team should be trained regarding US examination. Needless to say, with improving technology, consulting the US findings to an expert at a remote center for immediate medical advice is not impossible (Figure 1). As such, adding to the question(s) in the title, it would probably be high time to discuss where US should play: at the touch line or the corner area?



FIGURE 1. A simulation photograph illustrating how ultrasound examination can be performed during the match and be used simultaneously via telemedicine, as well.

Received: May 03, 2023

Accepted: May 09, 2023

Published online: May 18, 2023

Correspondence: Orhan Güvener, MD. Mersin Üniversitesi Tıp Fakültesi, Fiziksel ve Rehabilitasyon Tıp Anabilim Dalı, 33343 Yenişehir, Mersin, Türkiye.

E-mail: dr.orhanguvener@gmail.com

Doi: 10.52312/jdrs.2023.1201

Citation: Güvener O, Karanfil Y, Özçakar L. (How) Should ultrasound enter the match?. Jt Dis Relat Surg 2023;34(2):530. doi: 10.52312/jdrs.2023.1201

©2023 All right reserved by the Turkish Joint Diseases Foundation

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes (<http://creativecommons.org/licenses/by-nc/4.0/>).

Data Sharing Statement: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Author Contributions: All authors contributed equally to the article.

Conflict of Interest: The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

Funding: The authors received no financial support for the research and/or authorship of this article.

REFERENCES

- Özçakar L, Kara M, Chang KV, Çarlı AB, Akkaya N, Tok F, et al. Nineteen reasons why physiatrists should do musculoskeletal ultrasound: EURO-MUSCULUS/USPRM recommendations. Am J Phys Med Rehabil 2015;94:e45-9. doi: 10.1097/PHM.0000000000000223.
- Tok F, Özçakar L, De Muyneck M, Kara M, Vanderstraeten G. Musculoskeletal ultrasound for sports injuries. Eur J Phys Rehabil Med 2012;48:651-63.